



Voyxact[®]
 (sibeprenlimab-szsi)
 Injection 400 mg/2 mL

Exploring VOYXACT[®] with your care team

Fill out this discussion guide and bring it with you to your next appointment.

INDICATION

VOYXACT is a prescription medicine used to reduce protein in the urine (proteinuria) in adults with a kidney disease called primary immunoglobulin A nephropathy (IgAN) who are at risk for their disease getting worse. It is not known if VOYXACT is safe and effective in children.

This indication is approved under accelerated approval based on reduction of proteinuria. It has not been established whether VOYXACT slows kidney function decline over the long-term in patients with IgAN. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory clinical trial.

STEP 1: Get ready for your appointment

Review your latest labs.

Take a look at your last uPCR (urine protein-to-creatinine ratio). This test is important because it checks how much protein is in your urine.

Consider how you feel right now.

Are you experiencing symptoms like foamy urine, flank pain, swelling, or fatigue? Are you having side effects from your current medication(s)? Use the space below to describe any concerns you currently have. And make sure to let your nephrologist know about any new medical diagnoses or medications.

STEP 2: Start the conversation

Share your notes with your doctor about how you've been feeling. Use the space below to help discuss if VOYXACT could be an option for you.



REMEMBER:

It's important to remember that the lower your proteinuria levels, the better.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about VOYXACT?

- **Do not take VOYXACT if you** are allergic to any of the ingredients in VOYXACT. Stop using VOYXACT and get emergency help if you develop signs and symptoms of an allergic reaction, including:
 - breathing problems
 - dizziness or feel lightheaded
 - hives
 - swelling of the face, lips, mouth, tongue or throat
 - itching
 - skin rash

Please see Page 3 for Important Safety Information and QR codes linking to the full [Prescribing Information](#) and [Patient Information](#).

STEP 3: Ask follow-up questions

Characteristics of VOYXACT:

- How does VOYXACT work?
- How well does it work in lowering proteinuria?
- Will I need to change other parts of my IgAN care plan if I add VOYXACT?

Safety of VOYXACT:

- What are the most common side effects when taking VOYXACT? Is there anything I should watch out for?
- Are any additional labs or appointments needed when I start VOYXACT, beyond my usual monitoring?



VOYXACT is one self-administered injection, every 4 weeks, delivered to you in a prefilled syringe.

Other considerations:

- Am I eligible for VOYXACT?
- Are there any access and coverage resources available to patients starting VOYXACT?
- How do I get started on VOYXACT?
- How do I learn how to self-inject? Can someone else give me the injection?

IgAN=immunoglobulin A nephropathy.

STEP 4: Decide on next steps

Record any action items your doctor wants you to do before your next appointment.



If your doctor prescribes VOYXACT and you have questions about the medicine or need assistance, you can call Otsuka Patient Services at **833-VOYXACT** (869-9228), Monday through Friday, 8 AM–8 PM ET (except holidays).

IMPORTANT SAFETY INFORMATION (cont'd)

Before taking VOYXACT, tell your healthcare provider about all of your medical conditions, including if you:

- have an infection.
- have received or plan to receive therapy that can lower the ability of your immune system to fight infections (immunosuppressive therapy).
- have recently received or are scheduled to receive an immunization (vaccine). VOYXACT may affect your immune system response to vaccines and increase your risk of infections from live vaccines.
- are pregnant or plan to become pregnant. It is not known if VOYXACT will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if VOYXACT passes into your breast milk or if it will affect your baby. Talk to your healthcare provider about the best way to feed your baby during treatment with VOYXACT.

INDICATION and IMPORTANT SAFETY INFORMATION for VOYXACT[®] (sibeprenlimab-szsi)

INDICATION

VOYXACT is a prescription medicine used to reduce protein in the urine (proteinuria) in adults with a kidney disease called primary immunoglobulin A nephropathy (IgAN) who are at risk for their disease getting worse. It is not known if VOYXACT is safe and effective in children.

This indication is approved under accelerated approval based on reduction of proteinuria. It has not been established whether VOYXACT slows kidney function decline over the long-term in patients with IgAN. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory clinical trial.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about VOYXACT?

- **Do not take VOYXACT if you** are allergic to any of the ingredients in VOYXACT. Stop using VOYXACT and get emergency help if you develop signs and symptoms of an allergic reaction, including:
 - breathing problems
 - swelling of the face, lips, mouth, tongue or throat
 - dizziness or feel lightheaded
 - itching
 - hives
 - skin rash

Before taking VOYXACT, tell your healthcare provider about all of your medical conditions, including if you:

- have an infection.
- have received or plan to receive therapy that can lower the ability of your immune system to fight infections (immunosuppressive therapy).
- have recently received or are scheduled to receive an immunization (vaccine). VOYXACT may affect your immune system response to vaccines and increase your risk of infections from live vaccines.
- are pregnant or plan to become pregnant. It is not known if VOYXACT will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if VOYXACT passes into your breast milk or if it will affect your baby. Talk to your healthcare provider about the best way to feed your baby during treatment with VOYXACT.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

How should I take VOYXACT?

- **Read the detailed “Instructions for Use”** that come with VOYXACT for information about how to prepare and inject VOYXACT, and how to properly store and throw away (dispose of) used VOYXACT prefilled syringes.
- Use VOYXACT exactly as your healthcare provider tells you to.
- If you miss a dose of VOYXACT, inject the missed dose as soon as possible. Then take your next dose in 4 weeks.

What are the possible side effects of VOYXACT?


VOYXACT can cause serious side effects, including:

- **immunosuppression and increased risk of infections.** VOYXACT weakens your immune system, which may increase your risk of infections. Your healthcare provider will check you for signs and symptoms of infections before you start and during treatment with VOYXACT. Tell your healthcare provider right away if you develop any signs and symptoms of infection during treatment with VOYXACT, including:
 - fever
 - chills
 - feeling tired
 - cough
 - flu-like symptoms
 - aches and pain
 - warm, red, or painful skin


The most common side effects of VOYXACT include: upper respiratory tract infection and skin redness at your injection site.

These are not all of the possible side effects of VOYXACT.

Call your doctor for medical advice about side effects. You are encouraged to report side effects to FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.



Prescribing Information



Patient Information

This product information is intended for US consumers only.

VOYXACT is a registered trademark of Otsuka America Pharmaceutical, Inc.

© 2026 Otsuka America Pharmaceutical, Inc. All rights reserved.

January 2026

21US25EBC0170